



Rationale, Design and Early Trial performance of AOSpine North America Multi-Center Double Blind Randomized Controlled Trial of Safety and Efficacy of riluzole in CSM (CSM – Protect Trial).

Michael G. Fehlings MD PhD FRCS C FACS; Branko Kopjar MD

University of Toronto, Toronto, ON, Canada; University of Washington, Seattle, WA



Background

Cervical spondylotic myelopathy (CSM) is the most common cause of spinal cord impairment. While there is emerging evidence from the recently completed AOSpine North America prospective study that surgical decompression is an effective treatment for CSM, many patients have substantial residual neurological and functional impairment. Further improvement in treatment of CSM is warranted. Compelling evidence from preclinical models of nontraumatic and traumatic spinal cord injury (SCI) suggest a benefit of adding a neuroprotective drug which targets sodium/glutamate excitotoxicity to the treatment of patients with CSM undergoing surgical decompression.

Patients

A total of 270 (300 to adjust for loss-to-follow up) patients undergoing surgical decompression for CSM will be randomized in this ongoing prospective double-blinded controlled trial involving 15 sites in North America. Randomization will be 1:1 to riluzole 2x50mg daily for 14 days before the surgery and 28 days after the surgery or to the same regimen of placebo. Primary outcome measure is change in mJOA between baseline and 6 months following the surgery. Secondary outcomes include ASIA, SF36v2, NDI, EQ5D, Pain VAS and complications. Outcomes evaluations will occur at 6 and 12 months.

Statistical Design

Sample size of 270 subjects total will have 80% power to detect .35 Cohen's d effect size (i.e. 0.9 difference in mJOA). Study uses adaptive sequential design that allows sample size change during the interim analysis.

Results

Plan ID	Parameter
Type of the hypothesis	1-Sided
Type I Error (α)	0.025
Power ($1 - \beta$)	0.80
Randomization Ratio (Investigational vs. Control)	1:1
Planned Number of Interim Looks	2
Spacing of Looks	65%, 100%
Hypothesis to be Rejected	H0 or H1 (binding)
Boundary Family	Published Function
Boundary to Reject H0	O'Brien-Fleming
Boundary to Reject H1	Gamma (-2)
Difference of Means Assuming H ₁	0.9
Standard Deviation (σ)	2.57
Sample Size	270 (135 per arm)

Demographics		N (%)
Age (N = 147)		58.4 ± 10.12
Gender	Female	61 (41.22%)
	Male	87 (58.78%)
Race	White	124 (83.78%)
	African-American	11 (7.43%)
	Asian	6 (4.05%)
	American Indian or Alaska Native	1 (0.68%)
	Unknown	3 (2.03%)
	Other	3 (2.03%)
Surgery	Anterior	54 (42.19%)
	Posterior	72 (56.25%)
	Ant + Pos	2 (1.56%)

Primary and Secondary Outcomes

Outcome Measure	Mean (Standard Deviation)	
mJOA (N = 148)	11.8 (1.5)	
ASIA	Motor Total (N = 145)	95.3 (5.9)
	Sensory Light Touch (N = 143)	105.2 (11.5)
	Sensory Pin Prick (N = 138)	105.1 (10.6)
Pain VAS	Pain in Arm and Shoulder (N = 147)	5.4 (8.3)
	Pain in Neck (N = 147)	5.4 (8.3)
NDI (N = 145)	42.2 (20.0)	
Nurick (N=145)	3.2 (0.8)	

Conclusions

In spite of the benefits of the surgical intervention, patients with CSM experience significant residual impairment and neurological compromise. Adding neuroprotective treatment with riluzole may improve outcomes of surgery. This study will bring Level I evidence about efficacy of riluzole as adjuvant to surgical decompression in patients with moderate to severe CSM.

Acknowledgements

This study was sponsored by AOSpine North America Inc., a 501(c) 3 non-profit corporation.

